

NOTE

Apicoplast and Mitochondrion in Gametocytogenesis of *Plasmodium falciparum*^{▽†}

Noriko Okamoto,[‡] Timothy P. Spurck, Christopher D. Goodman, and Geoffrey I. McFadden*

Plant Cell Biology Research Centre, School of Botany, University of Melbourne, Parkville, Victoria 3010, Australia

Received 6 August 2008/Accepted 29 October 2008

Live cell imaging of human malaria parasites *Plasmodium falciparum* during gametocytogenesis revealed that the apicoplast does not grow, whereas the mitochondrion undergoes remarkable morphological development. A close connection of the two organelles is consistently maintained. The apicoplast and mitochondrion are not components of the male gametes, suggesting maternal inheritance.

The human malaria parasite *Plasmodium falciparum* is responsible for severe disease and persists as a major global health problem. *P. falciparum* belongs to the phylum *Apicomplexa* and harbors a relic plastid (apicoplast) acquired via secondary endosymbiosis (10, 13). Although the apicoplast has lost its photosynthetic ability (10), it still retains critical metabolic pathways such as type II fatty acid synthesis, isoprenoid synthesis, and part of heme synthesis (11, 20, 29, 39, 41). Because these pathways are fundamentally bacterial and distinct from equivalent human pathways, the apicoplast is a promising drug target (28).

Like all plastids, apicoplasts cannot be created de novo and must be vertically inherited. We previously demonstrated that both the apicoplast and the mitochondrion begin as small globular structures in the initial erythrocyte phase and then grow and branch before segmenting into multiple daughter organelles that are segregated into the daughter parasites (3, 15, 16, 20, 37, 39).

Despite detailed investigation of organelle inheritance in the asexual erythrocyte stages, little is known about other life cycle stages. The *falciparum* malaria parasite's life cycle is divided between the human and the mosquito. Transition from the human to the mosquito stages involves gametocytogenesis, which commences in erythrocytes and then proceeds in the mosquito gut, where the macrogametocyte (female) produces a single macrogamete and the microgametocyte (male) produces eight microgametes after three serial nuclear divisions. Gametes fuse to produce a zygote,

which eventually develops into motile forms ready for injection into a new human host.

We wanted to explore the behavior of the apicoplast and the mitochondrion in gametocytes (20, 27, 37, 39). We therefore created a new, tagged organelle line in the gametocyte-competent line 3D7 (2, 25, 30) with a novel fluorophore, mOrange (32), in the apicoplast for improved labeling. The 3D7 ACP(L):mOrange-CS(L):GFPmut2 line has red apicoplasts and green mitochondria, and examination of the asexual erythrocyte stages confirmed previous observations of organelle development and partitioning (37).

Five morphological developmental stages of gametocytes (14) are recognized (Fig. 1 and 2). In contrast to the apicoplast change during gametocytogenesis; the apicoplast stayed small and rounded or was only slightly elongated. Despite its morphological inertia in gametocytes, the apicoplast is functionally active. Sullivan et al. (35) demonstrated that inhibition of the apicoplast translation in the *P. bergii* gametocyte by thiostrepton blocks parasite transmission (12, 23). Furthermore, gametocyte transcriptomics shows upregulation of genes for apicoplast type II fatty acid biosynthesis (FASII), lipote synthase and the pyruvate dehydrogenase E1 alpha subunit (42), and at least two FASII enzymes are detected in gametocyte proteomes (18).

In contrast to the morphologically static apicoplast, the mitochondrion undergoes spectacular morphological development as gametocytes mature. In stage II gametocytes, the mitochondrion elongates and branches in a manner somewhat reminiscent of that in the asexual schizont cell, except that it forms a cluster around the small apicoplast (Fig. 1a and b; see also the movies in the supplemental material). The mitochondrion keeps this unusual association with the apicoplast throughout the whole gametocyte developmental process, typically appearing like a thicket of elongate mitochondrial profiles virtually encasing the apicoplast.

As the gametocyte elongates at Stage III, the mitochon-

* Corresponding author. Mailing address: School of Botany, University of Melbourne, Parkville 3010, Victoria, Australia. Phone: 61-414-189-905. Fax: 61-3-9347-1071. E-mail: gim@unimelb.edu.au.

† Supplemental material for this article may be found at <http://ec.asm.org/>.

‡ Present address: Department of Botany, University of British Columbia, Vancouver, BC V6T 1Z4, Canada.

[▽] Published ahead of print on 7 November 2008.




FIG. 1. Live cell images of 3D7 ACP(L):mOrange-CS(L):GFPmut2 gametocytes at stage II (a and b) and stage III (c to f), showing a small apicoplast (red) that is always encased by the mitochondrion (green). At stage III, the apicoplast is round (c and e) or slightly elongated (d and f). In contrast, the mitochondrion forms branches from the cluster that encase the apicoplast. Sometimes the GFP labeling for the mitochondrion is heterologous, but continuous (c and d). Scale bars, 3 μ m.

drion also elongates longitudinally (Fig. 1c to f; see also the movies in the supplemental material). In these cells, part of the mitochondrion is often folded a few times along the elongated apicoplast. As the gametocytes enlarge and mature (stage IV and V; Fig. 2), the mitochondrion in some cells forms a denser cluster with short, round branches that emerge from a center and encompass the apicoplast (Fig. 2a, c, and d; see also the movies in the supplemental material). However, the morphology is diverse and does not correspond to gametocyte sex or maturity.

Our observation of an expanding mitochondrion in the gametocyte is consistent with the activation of mitochondrial metabolism in gametocytes. Parasites in the asexual erythrocyte phase rely exclusively on cytosolic glycolysis and eschew the mitochondrial tricarboxylic acid (TCA) cycle. Indeed, Painter et al. suggest that mitochondrial electron

transport in erythrocytes is only necessary for the reduction of dihydroorotate dehydrogenase (4, 8, 27, 38). However, transcriptomics reveals that 15 of the 16 mitochondrial TCA cycle enzymes are upregulated in gametocytes (42), and nine of these enzymes are detected proteomically in females and four are detected in males (18). Gametocyte mitochondria also develop the tubular mitochondrial cristae that are largely absent in the asexual stages (21) but are required for full mitochondrial function. The elaboration of the mitochondria during gametocytogenesis reflects this development of mitochondrial activity.

The proximity of the mitochondrion and the apicoplast is intriguing. In the asexual phase, there is a constant physical association of the apicoplast and the mitochondrion (1, 15, 37), which even persists when the organelles are isolated (19, 26). This contact is proposed to facilitate metabolite




FIG. 2. Live cell images of 3D7 ACP(L):mOrange-CS(L):GFPmut2, at stage IV (a and b) and stage V (c and d), showing the apicoplast (red) and mitochondrion (green) association in macrogametocytes (a and c) and microgametocytes (b and d). The apicoplast (red) does not elongate or branch but stays small. Often, the mOrange signal for the apicoplast is weak in the mature microgametocyte (d). Scale bars, 3 μ m.

exchange (20, 39). Available evidence suggests that synthesis of heme by malaria parasites is a cooperative activity between the apicoplast and mitochondrion and that protoporphyrin intermediates are shuttled between the organelles (31, 39, 40). The gametocyte proteome includes at least one apicoplast and one mitochondrial enzyme for the putatively conjoined heme synthesis pathway (42), and the connection reported here may facilitate substrate shuttling. It has also been suggested that the two organelles may share tRNAs (16).

In the present study, we observed a single mitochondrion with the different segments visibly connected by thin strands. This is similar to rhodamine-123 staining of live gametocytes (17). The highly lobed mitochondrial structure may be responsible for the reports of multiple (four to eight) mitochondria based solely on electron microscopy of gametocytes of *P. falciparum* (33) and other *Plasmodium* species (21).

We also wanted to explore how the organelles might be inherited during sexual reproduction. In plants, both the plastid and the mitochondria are typically maternally inherited (22), but there are examples of biparental or paternal inheritance (24). We attempted to induce microgametogenesis (exflagellation) in our 3D7 ACP(L):mOrange-CS(L):GFPmut2 line without success. Instead, we used immunofluorescent markers for the mitochondrion and apicoplast to examine fixed parasites of the exflagellation-competent strain NF54 (25, 30).

We focused on the microgametogenesis, or exflagellation, since the obvious metamorphosis allows us to unambiguously distinguish mature gametes. NF54 parasites developed into gametocytes and the males could be induced to exflagellate as previously described (9). Seven minutes after the exflagellation induction, the microgametocyte begins to develop axonemes (Fig. 3a and b), retaining a single apicoplast and a single mitochondrion. At 30 min after the induction, when microgametes were being released, the single apicoplast and the single mitochondrion remain lodged in the original gametocyte cytoplasm and hence are not partitioned into the microgamete (Fig. 3c to f). The absence of mitochondria and apicoplasts in male gametes is consistent with genetic evidence for the maternal inheritance of the mitochondrial and apicoplast genomes (5), as well as electron microscopy (33, 34). The absence of apicoplasts and mitochondria in microgametes of other apicomplexan parasites (6, 7) suggests that this may be a common feature of the phylum (33).

Conclusions and future prospects. Gametocytogenesis in *P. falciparum* is the first step of the sexual reproduction, which is an essential bridging stage from humans into the mosquito host. In the present study, we made thorough morphological observations of the different gametocyte stages to demonstrate that the gametocyte has a single unelaborated apicoplast that undergoes minimal expansion through gametocytogenesis. In contrast, the single mitochondrion expands greatly to form a cluster around/along the apicoplast during




FIG. 3. Exflagellated microgametes. Immunofluorescence images of the mitochondrion in NF54 parasites, fixed 7 min (a and b) or 30 min (c to f) after exflagellation induction. (a to d) The mitochondrion (labeled with the anti-Hsp60 antibody [green]) stays single even after each microgamete (whose axoneme is labeled with the anti-alpha tubulin antibody [red]) is formed (a and b) and exflagellated (c and d). (e and f) The apicoplast (labeled with the anti-ACP antibody [green]) stays single and does not enter the exflagellated microgametes (whose axoneme is labeled with the anti-alpha tubulin antibody [red]). Scale bars, 3 μ m.

gametocyte development. The morphologies are congruent with metabolic activity in the two organelles at this stage.

Recent transcriptome and proteome analyses of the hepatocyte phase of the rodent malaria *P. yoelii* revealed high activity of the apicoplast and the mitochondrion (36), so it will be interesting to visualize the organelles in these stages, as well as sexual reproduction process in the mosquito host. A complete picture of what the organelles are doing at various life cycle phases will inform development of intervention strategies, particularly when and how to use drugs targeting apicoplast or mitochondrial metabolisms.

We thank to Alan F. Cowman (Walter and Eliza Hall Institute of Medical Research, Australia) for the NF54 strain and heat inactivated human serum. We also thank for Nirbhay Kumar (Johns Hopkins Malaria Research Institute) for advice on exflagellation. We thank the Australian Red Cross for human blood samples.

N.O. is funded by the Japan Society for Promotion of Sciences. G.M. is an ARC Federation Fellow and a Howard Hughes Medical Institute International Scholar. The project is supported by a Program Grant from the NH&MRC.

REFERENCES

1. Aikawa, M. 1966. The fine structure of the erythrocytic stages of three avian malarial parasites, *Plasmodium fallax*, *P. lophurae*, and *P. cathemerium*. *Am. J. Trop. Med. Hyg.* **15**:449–471.
2. Alano, P. 2007. *Plasmodium falciparum* gametocytes: still many secrets of a hidden life. *Mol. Microbiol.* **66**:291–302.
3. Bannister, L. H., J. M. Hopkins, R. E. Fowler, S. Krishna, and G. H. Mitchell. 2000. A brief illustrated guide to the ultrastructure of *Plasmodium falciparum* asexual blood stages. *Parasitol. Today* **16**:427–433.
4. Biagini, G. A., N. Fisher, N. Berry, P. A. Stocks, B. Meunier, D. P. Williams, R. Bonar-Law, P. G. Bray, A. Owen, P. M. O'Neill, and S. A. Ward. 2008. Acridinediones: selective and potent inhibitors of the malaria parasite mitochondrial bc1 complex. *Mol. Pharmacol.* **73**:1347–1355.
5. Creasey, A., K. Mendis, J. Carlton, D. Williamson, I. Wilson, and R. Carter. 1994. Maternal inheritance of extrachromosomal DNA in malaria parasites. *Mol. Biochem. Parasitol.* **65**:95–98.
6. Ferguson, D. J. P., S. A. Campbell, F. L. Henriquez, L. Phan, E. Mui, T. A. Richards, S. P. Muench, M. Allary, J. Z. Lu, S. T. Prigge, F. Tomley, M. W. Shirley, D. W. Rice, R. McLeod, and C. W. Roberts. 2007. Enzymes of type II fatty acid synthesis and apicoplast differentiation and division in *Eimeria tenella*. *Int. J. Parasitol.* **37**:33–51.
7. Ferguson, D. J. P., F. L. Henriquez, M. J. Kirisits, S. P. Muench, S. T. Prigge, D. W. Rice, C. W. Roberts, and R. L. McLeod. 2005. Maternal inheritance and stage-specific variation of the apicoplast in *Toxoplasma gondii* during development in the intermediate and definitive host. *Eukaryot. Cell* **4**:814–826.
8. Fisher, N., P. G. Bray, S. A. Ward, and G. A. Biagini. 2008. Malaria-parasite mitochondrial dehydrogenases as drug targets: too early to write the obituary. *Trends Parasitol.* **24**:9–10.
9. Fivelman, Q. L., L. McRobert, S. Sharp, C. J. Taylor, M. Saeed, C. A. Swales, C. J. Sutherland, and D. A. Baker. 2007. Improved synchronous production of *Plasmodium falciparum* gametocytes *in vitro*. *Mol. Biochem. Parasitol.* **154**:119–123.
10. Foth, B. J., and G. I. McFadden. 2003. The apicoplast: a plastid in *Plasmodium falciparum* and other apicomplexan parasites. *Int. Rev. Cytol.* **224**:57–110.
11. Goodman, C. D., and G. I. McFadden. 2007. Fatty acid biosynthesis as a drug target in apicomplexan parasites. *Curr. Drug Targets* **8**:15–30.
12. Goodman, C. D., V. Su, and G. I. McFadden. 2007. The effects of antibacterials on the malaria parasite *Plasmodium falciparum*. *Mol. Biochem. Parasitol.* **152**:181–191.
13. Gould, S. B., R. F. Waller, and G. I. McFadden. 2008. Plastid evolution. *Annu. Rev. Plant Biol.* **59**:491–517.
14. Hawking, F., M. E. Wilson, and K. Gammage. 1971. Evidence for cyclic development and short-lived maturity in gametocytes of *Plasmodium falciparum*. *Trans. R. Soc. Trop. Med. Hyg.* **65**:549–559.
15. Hopkins, J., R. Fowler, S. Krishna, I. Wilson, G. Mitchell, and L. Bannister. 1999. The plastid in *Plasmodium falciparum* asexual blood stages: a three-dimensional ultrastructural analysis. *Protist* **150**:283–295.
16. Howe, C. J., and S. Purton. 2007. The little genome of apicomplexan plastids: its raison d'être and a possible explanation for the 'delayed death' phenomenon. *Protist* **158**:121–133.
17. Kato, M., K. Tanabe, A. Miki, K. Ichimori, and S. Waki. 1990. Membrane potential of *Plasmodium falciparum* gametocytes monitored with rhodamine 123. *FEMS Microbiol. Lett.* **69**:283–288.
18. Khan, S. M., B. Franke-Fayard, G. R. Mair, E. Lasonder, C. J. Janse, M. Mann, and A. P. Waters. 2005. Proteome analysis of separated male and female gametocytes reveals novel sex-specific *Plasmodium* biology. *Cell* **121**:675–687.
19. Kobayashi, T., S. Sato, S. Takamiya, K. Komaki-Yasuda, K. Yano, A. Hirata, I. Onitsuka, M. Hata, F. Mi-ichi, T. Tanaka, T. Hase, A. Miyajima, S. Kawazu, Y. Watanabe, and K. Kita. 2007. Mitochondria and apicoplast of *Plasmodium falciparum*: behaviour on subcellular fractionation and the implication. *Mitochondrion* **7**:125–132.
20. Krungkrai, J. 2004. The multiple roles of the mitochondrion of the malarial parasite. *Parasitology* **129**:511–524.
21. Krungkrai, J., P. Prapunwattana, and S. R. Krungkrai. 2000. Ultrastructure and function of mitochondria in gametocytic stage of *Plasmodium falciparum*. *Parasite* **7**:19–26.
22. Kuroiwa, T., S. Kawano, S. Nishibayashi, and C. Sato. 1982. Epifluorescent microscopic evidence for maternal inheritance of chloroplast DNA. *Nature* **298**:481–483.
23. McConkey, G. A., M. J. Rogers, and T. F. McCutchan. 1997. Inhibition of *Plasmodium falciparum* protein synthesis: targeting the plastid-like organelle with thiostrepton. *J. Biol. Chem.* **272**:2046–2049.
24. Mogensen, H. L. 1996. The hows and whys of cytoplasmic inheritance in seed plants. *Am. J. Bot.* **83**:383–404.
25. Moore, J. M., N. Kumar, L. D. Shultz, and T. V. Rajan. 1995. Maintenance of the human malarial parasite, *Plasmodium falciparum*, in *scid* mice and transmission of gametocytes to mosquitos. *J. Exp. Med.* **181**:2265–2270.
26. Mullin, K. A., L. Lim, S. A. Ralph, T. P. Spurck, E. Handman, and G. I.

- McFadden. 2006. Membrane transporters in the relict plastid of malaria parasites. *Proc. Natl. Acad. Sci. USA* **103**:9572–9577.
27. Painter, H. J., J. M. Morrissey, M. W. Mather, and A. B. Vaidya. 2007. Specific role of mitochondrial electron transport in blood-stage *Plasmodium falciparum*. *Nature* **446**:88–91.
28. Ralph, S. A., M. C. D'Ombrain, and G. I. McFadden. 2001. The apicoplast as an antimalarial drug target. *Drug Resist. Update* **4**:145–151.
29. Ralph, S. A., G. G. van Dooren, R. F. Waller, M. J. Crawford, M. J. Fraunholz, B. J. Foth, C. J. Tonkin, D. S. Roos, and G. I. McFadden. 2004. Metabolic maps and functions of the *Plasmodium falciparum* apicoplast. *Nat. Rev. Microbiol.* **2**:203–216.
30. Rossario, V. 1981. Cloning of naturally occurring mixed infections of malaria parasites. *Science* **212**:1037–1038.
31. Sato, S., B. Clough, L. Coates, and R. J. M. (I.) Wilson. 2004. Enzymes for heme biosynthesis are found in both the mitochondrion and plastid of the malaria parasite *Plasmodium falciparum*. *Protist* **155**:117–125.
32. Shaner, N. C., R. E. Campbell, P. A. Steinbach, B. N. G. Giepmans, A. E. Palmer, and R. Y. Tsien. 2004. Improved monomeric red, orange and yellow fluorescent proteins derived from *Discosoma* sp. red fluorescent protein. *Nat. Biotechnol.* **22**:1567–1572.
33. Sinden, R. E., E. U. Canning, R. S. Bray, and M. E. Smalley. 1978. Gametocyte and gamete development in *Plasmodium falciparum*. *Proc. R. Soc. Lond. B* **201**:375–380.
34. Sinden, R. E., E. U. Canning, and B. Spain. 1976. Gametogenesis and fertilization in *Plasmodium yeolii nigeriensis*: transmission electron microscope study. *Proc. R. Soc. Lond. B* **193**:55–76.
35. Sullivan, M., J. Li, S. Kumar, M. J. Rogers, and T. F. McCutchan. 2000. Effects of interruption of apicoplast function on malaria infection, development, and transmission. *Mol. Biochem. Parasitol.* **109**:17–23.
36. Tarun, A. S., X. Peng, R. F. Dumpit, Y. Ogata, H. Silva-Rivera, N. Camargo, T. M. Daly, L. W. Bergman, and S. H. I. Kappe. 2008. A combined transcriptome and proteome survey of malaria parasite liver stages. *Proc. Natl. Acad. Sci. USA* **105**:305–310.
37. van Dooren, G. G., M. Marti, C. J. Tonkin, L. M. Stimmler, A. F. Cowman, and G. I. McFadden. 2005. Development of the endoplasmic reticulum, mitochondrion and apicoplast during the asexual life cycle of *Plasmodium falciparum*. *Mol. Microbiol.* **57**:405–419.
38. van Dooren, G. G., and G. I. McFadden. 2007. Malaria: differential parasite drive. *Nature* **450**:955–956.
39. van Dooren, G. G., L. M. Stimmler, and G. I. McFadden. 2006. Metabolic maps and functions of the *Plasmodium* mitochondrion. *FEMS Microbiol. Rev.* **30**:596–630.
40. Varadharajan, S., B. K. C. Sagar, P. N. Rangarajan, and G. Padmanaban. 2004. Localization of ferrochelatase in *Plasmodium falciparum*. *Biochem. J.* **384**:429–436.
41. Wiesner, J., and H. Jomaa. 2007. Isoprenoid biosynthesis of the apicoplast as drug target. *Curr. Drug Targets* **8**:3–13.
42. Young, J. A., Q. L. Fivelman, P. L. Blair, P. de la Vega, K. G. Le Roch, Y. Zhou, D. J. Carucci, D. A. Baker, and E. A. Winzeler. 2005. The *Plasmodium falciparum* sexual development transcriptome: a microarray analysis using ontology-based pattern identification. *Mol. Biochem. Parasitol.* **143**:67–79.